

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|--|
| (51) International Patent Classification⁶ : A61K 31/365 | A2 | (11) International Publication Number: WO 97/38689 (43) International Publication Date: 23 October 1997 (23.10.97) |
| (21) International Application Number: PCT/EP97/01800 (22) International Filing Date: 10 April 1997 (10.04.97) (30) Priority Data: 9607564.3 12 April 1996 (12.04.96) GB 9622028.0 24 October 1996 (24.10.96) GB (71) Applicant (for all designated States except US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): HAEBERLIN, Barbara [CH/CH]; Haselrain 77, CH-4125 Riehen (CH). MAK, Ching, Pong [GB/CH]; Ob dem Fichtenrain 20, CH-4106 Therwil (CH). MEINZER, Armin [DE/DE]; Fliederweg 2, D-79426 Buggingen (DE). VONDERSCHER, Jacky [FR/FR]; 29, rue de Bruebach, F-68400 Riedisheim (FR). (74) Agent: ROTH, Bernhard, M.; Novartis AG, Lichtstrasse 35, CH-4002 Basel (CH). | | (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i> |
| (54) Title: ENTERIC-COATED PHARMACEUTICAL COMPOSITIONS OF MYCOPHENOLATE (57) Abstract This invention provides a pharmaceutical composition comprising a mycophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract. | | |

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LJ | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

ENTERIC-COATED PHARMACEUTICAL COMPOSITIONS OF MYCOPHENOLATE

This invention relates to mycophenolic acid.

5 Mycophenolic acid, also referred to herein as MPA, was first isolated in 1896, and has been extensively investigated as a pharmaceutical of potential commercial interest. It is known to have anti-tumor, anti-viral, immunosuppressive, anti-psoriatic, and anti-inflammatory activity [see e.g. W.A. Lee et al, *Pharmaceutical Research* (1990), 7, p. 161 - 166 and references cited therein]. Publications have appeared on MPA as an anti-cancer agent by Lilly scientists, see e.g. M.J. Sweeney et al., *Cancer Research* (1972), 32, 1795-1802, and
10 by ICI scientists, see e.g. GB 1,157,099 and 1,203,328 and as an immunosuppressant agent see e.g. A. Mitsui et al. *J. Antibiotics* (1969) 22, p. 358-363. In the above-mentioned article by W.A. Lee et al it is stated that attempts have been made to increase the bio-availability or specificity of MPA by making derivatives. The poor bioavailability of the acid was thought to be caused by undetermined factors such as drug complexation in the gastro-
15 intestinal lumen, a narrow absorption window, metabolism before absorption etc.. The preparation of the morpholinoethyl ester, also known as mycophenolate mofetil (sometimes referred to herein as MMF), was described which had considerably higher bioavailability than MPA (100% for MMF and 43% for MPA). This derivative has been recently introduced commercially as an immunosuppressant for the treatment or prevention of organ
20 or tissue transplant rejection, at daily dosages of from about 200 mg to about 3 grams p.o., e.g. about 2 g p.o. Patient compliance with MMF is not ideal, inter alia, because of side-effects e.g. gastro-intestinal side effects, the origin of which is not known.

We have now found, after exhaustive testing, that mycophenolate salts when enteric coated or adapted to be released in the upper part of the intestines, e.g. in the duodenum, jejunum
25 and/or ileum, are effective, well-tolerated, pharmaceuticals particularly for immunosuppressive indications especially for the treatment or prevention of organ, tissue

or cellular allograft or xenograft rejection, e.g. after transplant, or the treatment or prevention of immune-mediated diseases (autoimmune diseases) and have interesting bioavailability and stability characteristics. Moreover fewer unit dosage forms are required to be administered than for MMF, leading to easier administration.

5 The present invention provides in one aspect a pharmaceutical composition comprising a myophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract (hereinafter referred to as a composition of the invention). The composition may be adapted in any conventional manner, preferably with means adapted to prevent release of the mycophenolate in the stomach and to ensure release in the upper
10 part of the intestinal tract. In a further aspect the invention provides a pharmaceutical composition comprising a coated pharmaceutically acceptable mycophenolate salt.

Such salts are cationic salts, e.g. of alkali metals, especially the sodium salts. Sodium mycophenolate salts are known, e.g. in South African Patent 68/4959. We prefer to use the mono-sodium salt. This may be obtained in crystalline form by recrystallization from
15 acetone/ethanol if necessary with water; Mpt. 189 - 191°C.

The invention provides, more specifically, a solid enteric-coated composition in unit dose form for oral application, the core of the composition containing sodium mycophenolate in solid or liquid form.

20 The term "core" comprises sodium mycophenolate (or other cationic salt) if desired in admixture with further physiologically acceptable material, that can be surrounded by an enteric-coating. The term "core" comprises, in a wide sense, not only tablets, pellets or granules but also capsules, e.g. soft or hard capsules of gelatine or starch. Such cores may be produced in conventional manner. We have found that the mycophenolate salts, particularly the sodium salt, are particularly interesting for the production of tablets. When
25 tablet cores are used they have preferably a hardness of from ca. 10 to 70 N.

The pellets or granules may, after application of the enteric-coating as described hereinafter may be used as such or to fill capsules, e.g. hard gelatine capsules. If desired the capsules may be alternatively enteric-coated, e.g. in conventional manner.

5 Other pharmaceutically acceptable ingredients may be present in the cores, e.g. those conventionally used in the preparation of pharmaceutically compositions, e.g. fillers, e.g. lactose, glidants, e.g. silica, and lubricants, e.g. magnesium stearate.

10 The term "enteric coating" comprises any pharmaceutically acceptable coating preventing the release of the active agent in the stomach and sufficiently disintegrating in the intestine tract (by contact with approximately neutral or alkaline intestine juices) to allow the resorption of the active agent through the walls of the intestinal tract. Various in vitro tests for determining whether or not a coating is classified as an enteric coating have been published in the pharmacopoeia of various countries.

15 More specifically, the term "enteric coating" as used herein refers to a coating which remains intact for at least 2 hours, in contact with artificial gastric juices such as HCl of pH 1 at 36 to 38°C and preferably thereafter disintegrates within 30 minutes in artificial intestinal juices such as a KH_2PO_4 buffered solution of pH 6.8.

The thickness of the coating may vary and depends inter alia on its permeability in water and acids. A typical coating may be about 16-30, e.g. 16-20 or to 25, mg on a size 1 gelatine capsule. Similar thicknesses may be applied in other formulations.

20 In general satisfactory results are obtained with a coating of 5 - 100 μm , preferably 20 - 80 μm thickness. The coating is suitably selected from macromolecular polymers. Suitable polymers are listed in e.g. L. Lachman et al. The Theory and Practice of Industrial Pharmacy, 3rd Ed, 1986, p. 365 - 373, H. Sucker et al, Pharmazeutische Technologie, Thieme, 1991, p. 355 - 359, Hagers Handbuch der pharmazeutischen Praxis, 4th Ed. Vol.
25 7, pages 739 to 742 and 766 to 778, (Springer Verlag, 1971) and Remington's Pharmaceutical Sciences, 13th Ed., pages 1689 to 1691 (Mack Publ., Co., 1970) and

comprise e.g. cellulose ester derivatives, cellulose ethers, acrylic resins, such as methylacrylate copolymers and copolymers of maleic acid and phthalic acid derivatives.

The preferred films are made from cellulose acetate phthalate and trimellitate; methacrylic acid copolymers, e.g. copolymers derived from methylacrylic acid and esters thereof, containing at least 40% methylacrylic acid; and especially hydroxypropyl methylcellulose phthalate.

Methylacrylates include those of molecular weight above 100,000 daltons based on, e.g. methylacrylate and methyl or ethyl methylacrylate in a ratio of about 1:1. Typical products include Endragit L, e.g. L 100-55, marketed by Rohm GmbH, Darmstadt, Germany.

10 Typical cellulose acetate phthalates have an acetyl content of 17-26% and a phthalate content of from 30-40% with a viscosity of ca. 45-90cP.

Typical cellulose acetate trimellitates have an acetyl content of 17-26%, a trimellityl content from 25 - 35 % with a viscosity of ca. 15-20 cS. An example of an appropriate cellulose acetate trimellitate is the marketed product CAT (Eastman Kodak Company, USA).

15 Hydroxypropyl methylcellulose phthalates, typically have a molecular weight of from 20,000 to 100,000 daltons e.g. 80,000 to 130,000 daltons, e.g. a hydroxypropyl content of from 5 to 10%, a methoxy content of from 18 to 24% and a phthalyl content from 21 to 35%.

20 An example of an appropriate cellulose acetate phthalate is the marketed product CAP (Eastman Kodak, Rochester N.Y., USA).

Examples of suitable hydroxypropyl methylcellulose phthalates are the marketed products having a hydroxypropyl content of from 6-10%, a methoxy content of from 20-24%, a phthalyl content of from 21-27%, a molecular weight of about 84,000 daltons known under the trade mark HP50 and available from Shin-Etsu Chemical Co. Ltd., Tokyo, Japan, and

having a hydroxypropyl content, a methoxy content, and a phthalyl content of 5-9%, 18-22% and 27-35% respectively, and a molecular weight of 78,000 daltons, known under the trademark HP55 and available from the same supplier.

A preferred coating is HP 50.

- 5 The enteric coating may be carried out in conventional manner, e.g. so that the cores are sprayed with a solution of the enteric-coating.

Suitable solvents for the enteric-coating are for example organic solvents, e.g. an alcohol such as ethanol, a ketone such as acetone, halogenated hydrocarbons such as CH_2Cl_2 or mixtures of such solvents, e.g. ethanol /acetone, e.g. 1:1 to 10:1.

- 10 Conveniently a softener such as di-n-butylphthalate or triacetin is added to such a solution, e.g. in a ratio of coating material to softener of from 1: about 0.05 to about 0.3.

If desired for cellulose phthalates and other acidic coating materials an ammonium salt may be found and an aqueous solution may be used.

A fluidized bed coater may be used for coating.

- 15 Conveniently the cores are treated at room temperature or warmed up to 40°C e.g. by means of warm air of 40° up to 70°C, before spraying. To avoid a sticking of the cores the spray procedure is preferably interrupted at certain time intervals and the cores then warmed up again. It is, however, also possible to proceed without interruption of the spray procedure, e.g. by automatic regulation of the spray amount taking into account the temperature of
20 exhaust air and/or cores.

The spray pressure may vary within wide ranges, in general satisfactory results are obtained with a spray pressure of from about 1 to about 1.5 bar.

The compositions of the invention are useful as immunosuppressants as indicated by standard tests.

The activity and characteristics of the compositions of the invention may be indicated in standard

- 5 a) clinical trials, e.g. observing the first acute rejection episodes or treatment failure six months after transplant of kidneys or maintaining a rejection - free state within 6 months after initiation of treatment with the invention. The compositions of the invention are administered at a dose in the range of 0.5 to 2.0 g/day e.g. about 1.5 g /day and decrease the acute rejection rates when administered during the period
10 around transplant surgery, and maintain a rejection-free state in patients who are 3 months or more after transplantation. Thus the compositions of the invention may be administered during the initial 72 hours after transplantation at dose of about 0.5 g administered twice a day in combination with a conventional steroid and cyclosporin, e.g. as NEORAL for which the cyclosporin dose is the conventional dose e.g. ca. 8 ± 3 mg/kg for renal transplants. The steroid dose is to be
15 administered at about 2.5 mg /kg for 4 days after transplant, 1 mg/kg thereafter for 1 week, 0.6 mg/kg thereafter for 2 weeks thereafter 0.3 mg/kg for 1 month for prednisone.

and in

- 20 b) animal trials e.g. observing the kidney allograft reaction in rat. In this test one kidney from a female fisher 344 rat is transplanted onto the renal vessel of a unilaterally (left side) nephrectomized WF recipient rat using an end-to-end anastomosis. Ureteric anastomosis is also end-to-end. Treatment commences on the day of transplantation and is continued for 14 days. A contralateral nephrectomy is
25 done seven days after transplantation, leaving the recipient relying on the performance of the donor kidney. Survival of the graft recipient is taken as the parameter for a functional graft. Typical doses of the compositions of the invention

are from about 1 to 30 mg/kg p.o.

The compositions of the invention are particularly useful for the following conditions:

- 5 a) Treatment and prevention of native or transgenic organ, tissue or cellular allograft or xenograft transplant rejection, e.g. for the treatment of recipients of e.g. heart, lung, combined heart-lung, liver, kidney, pancreatic, skin, pancreatic islet cell, neural cell or corneal transplant; including treatment and prevention of acute rejection; treatment and prevention of hyperacute rejection, e.g. as associated with xenograft rejection; and treatment and prevention of chronic rejection, e.g. as associated with graft-vessel disease. The compositions of the invention are also indicated for the treatment and
- 10 prevention of graft-versus-host disease, such as following bone marrow transplantation.
- b) Treatment and prevention of autoimmune diseases, e.g. immune-mediated diseases and inflammatory conditions, in particular inflammatory conditions with an etiology including an immunological component such as arthritis (for example rheumatoid arthritis, arthritis chronica progrediente and arthritis deformans) and rheumatic
- 15 diseases. Specific immune-mediated diseases for which the compositions of the invention may be employed include, autoimmune hematological disorders, including, but not limited to hemolytic anaemia, aplastic anaemia, pure red cell anaemia and idiopathic thrombocytopenia), systemic lupus erythematosus, polychondritis, sclerodoma, Wegener granulosis, dermatomyositis, polymyositis, chronic active
- 20 hepatitis, primary biliary cirrhosis, myasthenia gravis, psoriasis, Steven-Johnson syndrome, pemphigus, idiopathic sprue, inflammatory bowel diseases (including e.g. ulcerative colitis and Crohn's disease), endocrine ophthalmopathy, Graves disease, sarcoidosis, multiple sclerosis, juvenile diabetes (diabetes mellitus type I), non-infectious uveitis (anterior and posterior), keratoconjunctivitis sicca and vernal
- 25 keratoconjunctivitis, interstitial lung fibrosis, psoriatic arthritis, vasculitis, glomerulonephritides (with and without nephrotic syndrome, e.g. including idiopathic nephrotic syndrome or minimal change nephropathy) and juvenile dermatomyositis.

Appropriate dosages of the compositions of the invention will of course vary, e.g. depending on the condition to be treated (for example the disease type or the nature of resistance), the MPA salt used, the effect desired and the mode of administration.

5 In general however satisfactory results are obtained on administration e.g. orally at dosages on the order of from about 1 to about 30 mg salt per kg animal body weight per day, administered once or in divided doses up to 4 times per day. Suitable daily dosages for patients are thus in the order of 200 mg to 3 g p.o. salt e.g. from about 50 to 100% that of mycophenolate mofetil. For the preferred mono sodium salt the dosage of the salt is about two thirds that of mycophenolate mofetil.

10 Representative unit dosage forms contain from about 50 mg, e.g. 100 mg, to about 1.5 g of the pharmaceutically acceptable mycophenolate salt.

The bioavailability characteristics of compositions of the invention may be determined in conventional manner, e.g. by oral administration to beagle dogs. Dosages are typically 50 mg salt animal e.g. ca 3- 5 mg salt /kg animal body weight. Dogs are adult (ca. 10 kg e.g.
15 6- 14 kg) and fasted. Three hours after administration ca. 200 g food is administered. Blood samples are taken from the cephalic vein, before administration and 10, 30, and 45 minutes, 1, 1.5, 2, 3, 4, 6, 8, 12, and 24 hours, after administration. Plasma levels of free MPA are determined by HPLC analysis (with UV detection).

20 In a relative bioavailability trial as described above in male beagle dogs dosages of 3.8 mg salt/kg animal body weight p.o. were administered with the Example 1 composition as described hereinafter and with a MPA or MMF formulation corresponding to the Example 1 composition but containing an identical amount of MPA or commercially available MMF.

Results are as follows:-

| | MPA (AUC Relative Bioavailability, Frel [ng.hr.ml ⁻¹]) | Ex 1 | MPA | MMF |
|----|--|------------|------------|------------|
| 5 | Mean | 4612 (218) | 3579 (174) | 2709 (100) |
| | Median | 4204 (168) | 2911 (182) | 2513 (100) |
| | SD | 939 | 1889 | 1363 |
| | CV | 20 | 53 | 50 |
| | Cmax [ng/ml] (Relative Cmax) | | | |
| 10 | Mean | 5391 (313) | 3683 (227) | 2052 (100) |
| | Median | 5359 (367) | 2719 (172) | 1462 (100) |
| | SD | 1847 | 2504 | 945 |
| | CV (%) | 34 (46) | 68 (87) | 46 (0) |

15 The coefficients of variation (CV) of AUC (20%) and Cmax (34%) of the Example 1 composition are significantly less than those of the reference compositions, indicating less inter-subject and intra-subject variability with the Example 1 composition.

The area under the curve (AUC) and Cmax with the Example 1 composition are higher than those of the reference compositions.

20 Naturally the advantageous bioavailability characteristics of the present compositions may be ascertained in standard clinical bioavailability trials. For example, doses from 200 mg to 1.5 g of the Example 1 composition and MPA, and MMF may be administered to 12 healthy volunteers in single doses in a cross-over trial. Increased AUC and C_{max} may be observed for the Example 1 composition.

25 The compositions of the present invention are surprisingly tolerated better than MMF, inducing less gastro-intestinal side effects such as diarrhoea and burning. They show less

long term side effects e.g. in the colon.

The compositions of the invention may be administered as the sole active ingredient or with another immunosuppressant e.g. together with simultaneous or separate administration of other immunosuppressants, for example, in immunosuppressive applications such as prevention and treatment of graft vs. host disease, transplant rejection, or immune-mediated disease, the compositions of the invention may be used in combination with cyclosporins or ascomycins, or their immunosuppressive analogs, e.g. cyclosporin A, FK- 506 (tacrolimus), etc., rapamycin; corticosteroids; cyclophosphamide; azathioprine; methotrexate; brequinar; leflunomide; mizoribine; deoxyspergualin; analogues thereof, and immunosuppressive monoclonal antibodies, e.g., monoclonal antibodies to leukocyte receptors, e.g. MHC, CD2, CD3, CD4, CD7, CD25, CD28, CTLA4, B7, CD45, or CS58 or their ligands; or other immunomodulatory compounds.

When the compositions of the invention are co-administered with such other immunosuppressants the dosages of the other immunosuppressants may be reduced e.g. to one-half to one-third their dosages when used alone.

Representative doses for ciclosporin to be used are e.g. 1 to 10, e.g. 1 to 2 mg/kg/day.

The present invention provides in another aspect the use, method and compositions as defined hereinafter in the claims.

Insofar as details of excipients are not described herein, these are known, or available e.g. in the Handbook of Pharmaceutical Excipients, Second Edition, edited by Ainley Wade and Paul J. Weller, American Pharmaceutical Association, Washington, USA and Pharmaceutical Press, London; and Lexikon der Hilfsstoffe für Pharmazie, Kosmetik and angrenzende Gebiete edited by H.P. Fiedler, 4th Edition, Editio Cantor, Aulendorf and earlier editions.

Following is a description by way of example only of compositions of this invention:

EXAMPLE 1:COMPOSITIONCapsule contents

| | | |
|---|---------------------------------------|------------------------|
| | MPA mono sodium salt | 53.43 mg (= 50 mg MPA) |
| 5 | Lactose (1:1 mixture of 100/200 mesh) | 256.57 mg |
| | Silica (Aerosil) | 3.1 mg |
| | Magnesium stearate | <u>1.55 mg</u> |
| | | <u>314.65 mg</u> |

Capsule is size 1

10 Enteric coating (ca 17 mg)

Hydroxypropyl methyl cellulose phthalate (HP50) 9 parts

Triacetin 1 part

Procedure

15 The capsule ingredients are mixed and filled into size 1 capsules. The capsules are coated in a fluidized bed coater with a solution of the enteric coating ingredients in ethanol (containing 10% acetone). The coating on each capsule is about 17 mg. The capsules meet the enteric coating test described herein and do not disintegrate within 2 hours in artificial gastric juices (pH 1, HCl). The compositions are stable, e.g for 2 years at room temperature.

20 If desired larger capsules containing 534.3 mg MPA mono sodium salt may be made in analogous manner, reducing the amount of lactose. These are well tolerated in clinical trials.

EXAMPLE 2:

Capsules of size 1 are made up as in Example 1. A solution for enteric coating is made up as follows:

| | | |
|----|---|--------|
| | Hydroxypropyl methyl cellulose phthalate (HP50) | 270 g |
| 25 | Triacetin | 30 g |
| | Acetone | 900 g |
| | Ethanol | 1800 g |

600 g of this enteric coating solution are used for 1 kg of capsules (ca. 2400). The amount of coating applied to each capsule is about 25 mg giving a film thickness of 5-6 mg/cm².

CLAIMS

1. A pharmaceutical composition comprising a mycophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract.
- 5 2. A pharmaceutical composition comprising an enteric coated pharmaceutically acceptable mycophenolate salt.
3. Use of an enteric coated pharmaceutically acceptable mycophenolate salt or composition of claim 1 or 2 in the preparation of a medicament for immunosuppression, particularly for prevention or treatment of native or transgenic organ, tissue or cellular allograft or xenograft transplant rejection; for the treatment or
10 prevention of immune-mediated and/or inflammatory diseases; optionally with the simultaneous or separate administration of another immunosuppressant.
4. A method of immunosuppressing a subject which comprises administering a therapeutically effective amount of enteric coated pharmaceutically acceptable mycophenolate or a composition of claim 1 or 2 to a subject in need of such
15 immunosuppression, optionally with the simultaneous or separate administration of another immunosuppressant.
5. A composition containing an enteric coated pharmaceutically acceptable mycophenolate salt or a composition of claim 1 or 2 and another immunosuppressant for simultaneous, sequential or separate administration.
- 20 6. A composition, use or method according to any preceding claim wherein the salt is the mono-sodium salt.
7. A composition, use or method according to any preceding claim wherein another immunosuppressant is present, e.g. ciclosporin.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|---|--|
| (51) International Patent Classification ⁶: A61K 31/365 | A3 | (11) International Publication Number: WO 97/38689 (43) International Publication Date: 23 October 1997 (23.10.97) |
| (21) International Application Number: PCT/EP97/01800 (22) International Filing Date: 10 April 1997 (10.04.97) (30) Priority Data: 9607564.3 12 April 1996 (12.04.96) GB 9622028.0 24 October 1996 (24.10.96) GB (71) Applicant (for all designated States except US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): HAEBERLIN, Barbara [CH/CH]; Haselrain 77, CH-4125 Riehen (CH); MAK, Ching, Pong [GB/CH]; Ob dem Fichtenrain 20, CH-4106 Therwil (CH); MEINZER, Armin [DE/DE]; Fliederweg 2, D-79426 Buggingen (DE); VONDERSCHER, Jacky [FR/FR]; 29, rue de Bruebach, F-68400 Riedisheim (FR). (74) Agent: ROTH, Bernhard, M.; Novartis AG, Lichtstrasse 35, CH-4002 Basel (CH). | (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 4 December 1997 (04.12.97) | |
| (54) Title: ENTERIC-COATED PHARMACEUTICAL COMPOSITIONS OF MYCOPHENOLATE (57) Abstract <p>This invention provides a pharmaceutical composition comprising a mycophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract.</p> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | | | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakhstan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/EP 97/01800

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/365

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| X | GB 1 203 328 A (ICI LIMITED) 26 August 1970 cited in the application see page 3; example 1 see page 1, line 21 --- | 1,5 |
| X | BE 704 326 A (ICI LIMITED) 26 March 1968 see page 4; example 1 see claims 1,2 --- | 1 |
| X | US 3 705 946 A (DYKE RICHARD W ET AL) 12 December 1972 see column 2, line 62 - line 70 --- | 1,6 |
| X | WO 94 26266 A (SYNTEX INC) 24 November 1994 see page 4, line 36 - page 5, line 26 --- | 1 |
| | --- -/-- | |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

9 October 1997

Date of mailing of the international search report

28.10.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 eponi,
Fax: (+31-70) 340-3016

Authorized officer

Boulois, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/01800

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| A | MARINARI R. ET AL: "Mycophenolic acid in the treatment of psoriasis" ARCH. DERMATOL., vol. 113, 1977, pages 930-932, XP002043018 see page 932; table 2 --- | 1-7 |
| A | GRINYO J. ET AL: "Placebo controlled study of mycophenolate mofetil combined with cyclosporin and corticosteroids for prevention of acute rejection" LANCET, vol. 345, 1995, pages 1321-1325, XP002043019 see page 1321, paragraph 1 --- | 3-5,7 |
| A | SUCKER H. ET AL: "Pharmazeutische Technologie" 1991, THIEME VERLAG, STUTTGART XP002043101 189560 cited in the application see page 356 - page 361 ----- | 1-7 |

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 97/01800

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 4,6,7
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.